



US005202253A

United States Patent [19]

Esmon et al.

[11] **Patent Number:** **5,202,253**[45] **Date of Patent:** **Apr. 13, 1993**

[54] **MONOCLONAL ANTIBODY SPECIFIC FOR PROTEIN C AND ANTIBODY PURIFICATION METHOD**

[75] **Inventors:** Charles T. Esmon; Naomi L. Esmon, both of Oklahoma City, Okla.

[73] **Assignee:** Oklahoma Medical Research Foundation, Oklahoma City, Okla.

[21] **Appl. No.:** 730,040

[22] **Filed:** Jul. 12, 1991

Related U.S. Application Data

[63] Continuation of Ser. No. 292,447, Dec. 30, 1988, abandoned.

[51] **Int. Cl.⁵** C12N 5/20; C12N 15/02; C07K 15/28; C07K 3/20

[52] **U.S. Cl.** 435/240.27; 530/388.25; 530/413; 435/172.2; 435/70.21

[58] **Field of Search** 530/388.25, 413; 435/240.27, 172.2, 70.21

[56] **References Cited**

U.S. PATENT DOCUMENTS

4,506,009 3/1985 Lenhoff et al. 435/7.5
5,147,638 9/1992 Esmon et al. 424/85.8

OTHER PUBLICATIONS

Seligsohn, et al., *The New England Journal of Medicine*, vol. 310 No. 9, pp. 559-462 (Mar. 1, 1984).

Vukovich, et al., *British Journal of Haematology* 70, pp. 435-440 (1988).

C. T. Esmon, et al., Joint IABS/CSL Symposium on Standardization in Blood Fractionation including Coagulation Factors, Melbourne, Australia, 1986 *Develop. Biol. Standard*, vol. 67, pp. 51-57 (S. Karger, Basel, 1987).

C. T. Esmon, et al., *Journ. of Biol. Chem.* vol. 264, No. 9 pp. 4743-4746 (Mar. 25, 1989).

C. T. Esmon, *Science* vol. 235, pp. 1348-1352 (Mar. 13, 1987).

A. K. Ohlin, et al., *Journ. of Biol. Chem.* vol. 262, No. 28 pp. 13798-13804 (Oct. 5, 1987).

Laurell, et al., *FEBS Letters* vol. 191, No. 1, pp. 75-81 (Oct. 1985).

Ohlin, et al., *Thrombosis and Haemostasis* 58(1): 230 (Jul. 1987).

Freyssinet, et al., *Thrombosis and Haemostasis* 58(1):230 (Jul., 1987).

Suzuki, et al., *J. Biochem.* 97, 127-138 (1985).

Stearns, et al., *Journ. of Biol. Chem.*, vol. 263, No. 2, pp. 826-832 (Jan. 15, 1988).

Beckmann, et al., *Nucl. Acids Res.* vol. 13, No. 14 pp. 5233-5247 (1985).

Sugo et al. *Thromb. Hemost.* 58(1) 229 1987.

Esmon et al., *Develop Biol. Standard.*, 67: 75-82, 1987.

Dreyfus et al., *New England Journal of Medicine* 325: 1563-1568, 1991.

Taylor et al. *J. Clin. Invest.* 79: 918-25 1987.

Ikeda et al. *Thromb. Res* 39: 297-306 1985.

Goding "Monoclonal Antibodies" Academic Press, 1983 pp. 111-113.

Primary Examiner—David L. Lacey

Assistant Examiner—Paula Hutzell

Attorney, Agent, or Firm—Kilpatrick & Cody

[57]

ABSTRACT

A Ca²⁺ dependent monoclonal antibody that specifically binds to a specific twelve peptide sequence (E D Q V D P R L I D G K) in the activation region of the Protein C. The antibody does not bind to Activated Protein C and can be used to inhibit activation of Protein C by thrombin-thrombomodulin. The antibody can be isolated from cell culture or ascites fluid in large quantities by affinity chromatography with mild conditions using the peptide bound to an immobilized substrate. The antibody has a number of specific uses in isolation and characterization of Protein C and as a model for the design of Ca²⁺ dependent antibodies for the isolation of other proteins, as a diagnostic, and as a therapeutic to prevent activation of Protein C. The Protein C can be naturally produced or produced by expression of the recombinant gene. Advantages of the antibody in purification of Protein C include the specificity for Protein C and not Activated Protein C, and the unique Ca²⁺-peptide binding specificity which allows the binding site to be protected when it is being immobilized on the chromatographic support. In vivo, the antibody has been demonstrated to inhibit tumor growth. The antibody can also be used to promote clotting in patients having high levels of Factor VIII inhibitors.

5 Claims, 2 Drawing Sheets